ChemRisk Document No. 1300

SECRET COVER SHEET

FV.12 2/20/13

Acid Recov

DAK RIDGE NATIONAL LABORATORY Operated By

CARBIDE AND CARBON CHEMICALS COMPANY

ш

POST OFFICE BOX P OAK RIDGE, TENNESSEE

ORNL CENTRAL FILES NUMBER

52-10-_

DATE:

October 8, 1952

COPY NO. 2-H

SUBJECT:

ICDINE IN THE ACID RECOVERY VENT SCRUBBER

TO:

R. E. Tomlinson

FROM:

J. W. Landry

DISTRIBUTION:

1. R. E. Tomlinson

2. F. L. Steahly

3. H. K. Jackson J. W. Landry

Chasification Cancelled

This document has been approved for release to the public by:


CLASSIFICATION CANCELLED

Date

Single rereview of CCRP-declassified documents was authorized by DOE Office of Declassification memo of August 22, 1994.

This document of its contents in any manne an unauthorized person is prohibited.

COVER SHEET



OAK RIDGE NATIONAL LABORATORY

OPERATED BY

CARBIDE AND CARBON CHEMICALS CORPORATION

FOR THE

ATOMIC ENERGY COMMISSION

POST OFFICE BOX P

OAK RIDGE, TENNESSEE

October 8, 1952

CLASSIFICATION CANCELLED

Of Morgan 1-31-95

Dsimature

Date

Single rereview of CCRP-declassified documents was authorized by DOE Office of freclassification memo of August 22, 1994

Mr. R. E. Tomlinson

Hanford Engineering Works

Richland, Washington

Subject: IODINE IN THE ACID RECOVERY VENT SCRUBBER

The analysis of the samples of 50% NaOH solution drawn October 3 from the acid recovery vent scrubber has been received:


Gross beta Iodine Iodine calculated to	10-3-52 10-6-52 10-3-52	6.27x10 ¹ beta counts/min/ml* 2.70x10 ¹ beta counts/min/ml 3.48x10 ¹ beta counts/min/ml
Ruthenium Zirconium	9-11-52 10-4-52 10-6-52	6.91x10 ⁴ beta counts/min/ml 3.28x10 ³ beta counts/min/ml 5.88x10 ³ beta counts/min/ml
Niobium Total Rare Earths	10-4-52	1.31x10 ³ beta counts/min/ml 8.38x10 ³ beta counts/min/ml


^{*} counted at 10.18% geometry

The iodine was present from thirty-six hours operation of acidrecovery during run HCP-14** which was made from sixty-five day cooled (to 9-11-52) 465 NWD/T material, the iodine in the four previous runs being negligible:

HCP-13	342 day cooled
HCP-12	310 day cooled
HCP-11	114 day cooled
HCP-10	126 day cooled

** HCP-14 EAW: Iodine = 0.03% of gross beta activity. The IAW was 9.14x10 gross beta counts/min/ml and 2.7x106 iodine beta counts/min/ml.





A leak of 50% of more of the stream at "L" developed in the beginning of the run and resulted in termination after thirty-six hours.

The average proportions of TRE, Ru, Zr, and No in the EAW for the five runs were approximately:

TRE	65% o	of gross	beta	activity
Ru	8% c	of gross	beta	activity
Zr				activity
Nb				activity

Instrument air exceeded the scrubber capacity on several occasions and scrubber efficiency is therefore questionable. The HCP-14 MAW boiled with an estimated acidity of 10.5 N HMO, and the preponderance of iodine in the scrubber sample seems sufficient to show iodine volatilized. The anomalous proportions between the other scrubber contaminants might indicate some preferential carryover via volatilization but not enough is known of the scrubber efficiency and TRE analysis for definite conclusions.

Very truly yours,

OAK RIDGE NATIONAL LABORATORY

J. W. Landry Chemical Technology Division

And the second s

JWL: ach

